

DCT Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9956A

Product Information

Application WB, IHC-P, E **Primary Accession** P40126

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalIsotypeRabbit IgGClone NamesRB23479Calculated MW59145Antigen Region62-89

Additional Information

Gene ID 1638

Other Names L-dopachrome tautomerase, DCT, DT, L-dopachrome Delta-isomerase,

Tyrosinase-related protein 2, TRP-2, TRP2, DCT, TYRP2

Target/Specificity This DCT antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 62-89 amino acids from the N-terminal

region of human DCT.

Dilution WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions DCT Antibody (N-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name DCT (HGNC:2709)

Synonyms TYRP2

Function Plays a role in melanin biosynthesis (PubMed: <u>33100333</u>). Catalyzes the

conversion of L-dopachrome into 5,6-dihydroxyindole-2- carboxylic acid

(DHICA).


Cellular Location

Melanosome membrane; Single-pass type I membrane protein. Melanosome {ECO:0000250 | UniProtKB:P29812}. Note=Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 {ECO:0000250 | UniProtKB:P29812}


References

Edwards, M., et al. PLoS Genet. 6 (3), E1000867 (2010) Khong, H.T., et al. J. Immunol. 168(2):951-956(2002) Sugimoto, H., et al. Biochemistry 38(11):3268-3279(1999)

Images



DCT Antibody (N-term) (Cat. #AP9956a) IHC analysis in formalin fixed and paraffin embedded breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the DCT Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Western blot analysis of DCT Antibody (N-term) (Cat. #AP9956a) in A375 cell line lysates (35ug/lane). DCT (arrow) was detected using the purified Pab.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.