

VAP1 Rabbit pAb

Catalog # AP99596

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q16853
Other Accession	Q16853
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84622
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human VAP1
Epitope Specificity	201-300/763
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the copper/topaquinone oxidase family.
Post-translational modifications	Topaquinone (TPQ) is generated by copper-dependent autoxidation of a specific tyrosyl residue. N- and O-glycosylated.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the semicarbazide-sensitive amine oxidase family. Copper amine oxidases catalyze the oxidative conversion of amines to aldehydes in the presence of copper and quinone cofactor. The encoded protein is localized to the cell surface, has adhesive properties as well as monoamine oxidase activity, and may be involved in leukocyte trafficking. Alterations in levels of the encoded protein may be associated with many diseases, including diabetes mellitus. A pseudogene of this gene has been described and is located approximately 9-kb downstream on the same chromosome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2013]

Additional Information

Gene ID	8639
Other Names	HPAO; SSAO; VAP-1; VAP1; AOC3_HUMAN; AOC3; Amine oxidase copper-containing 3; Copper amine oxidase; Semicarbazide-sensitive amine oxidase (SSAO); Vascular adhesion protein 1 (VAP-1); 1.4.3.21; AOC3_RAT; VP97; amine oxidase copper containing 3; vascular adhesion protein 1
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	AOC3 (HGNC:550)
Synonyms	VAP1
Function	Catalyzes the oxidative deamination of primary amines to the corresponding aldehydes with the concomitant production of hydrogen peroxide and ammonia (PubMed: 19588076 , PubMed: 24304424 , PubMed: 9653080). Has a preference for the primary monoamines methylamine and benzylamine (PubMed: 19588076 , PubMed: 9653080). Could also act on 2-phenylethylamine but much less efficiently (PubMed: 19588076). At endothelial cells surface can also function as a cell adhesion protein that participates in lymphocyte extravasation and recirculation by mediating the binding of lymphocytes to peripheral lymph node vascular endothelial cells in an L-selectin-independent fashion (PubMed: 9254657 , PubMed: 9653080).
Cellular Location	Cell membrane; Single-pass type II membrane protein
Tissue Location	Strongly expressed on the high endothelial venules of peripheral lymph nodes and on hepatic endothelia. Also highly expressed in appendix, lung and small intestine. Expressed also in adipose tissue, in bone marrow, colon, heart, kidney, ovary, pancreas, placenta, prostate, skeletal muscle, spleen and testis. Isoform 2 seems to be the predominant transcript in fetal kidneys, fetal cartilage and fetal tonsils. The highest relative expression of isoform 2 occurs in skeletal muscle, heart, pancreas, kidney, and lung

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.