

# DLL4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9964a

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">Q9NR61</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB23696
<b>Calculated MW</b>	74605
<b>Antigen Region</b>	625-652

## Additional Information

---

<b>Gene ID</b>	54567
<b>Other Names</b>	Delta-like protein 4, Drosophila Delta homolog 4, Delta4, DLL4
<b>Target/Specificity</b>	This DLL4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 625-652 amino acids from the C-terminal region of human DLL4.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	DLL4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	DLL4
<b>Function</b>	Involved in the Notch signaling pathway as Notch ligand (PubMed: <a href="#">11134954</a> ). Activates NOTCH1 and NOTCH4. Involved in angiogenesis; negatively regulates endothelial cell proliferation and migration and angiogenic sprouting (PubMed: <a href="#">20616313</a> ). Essential for retinal progenitor

proliferation. Required for suppressing rod fates in late retinal progenitors as well as for proper generation of other retinal cell types (By similarity). During spinal cord neurogenesis, inhibits V2a interneuron fate (PubMed:[17728344](#)).

<b>Cellular Location</b>	Cell membrane; Single-pass type I membrane protein
<b>Tissue Location</b>	Expressed in vascular endothelium.

## Background

---

DLL4 is a homolog of the Drosophila delta gene. The delta gene family encodes Notch ligands that are characterized by a DSL domain, EGF repeats, and a transmembrane domain.


## References

---

Emuss, V., et al. PLoS Pathog. 5 (10), E1000616 (2009)  
Ferrari-Toninelli, G., et al. Dev Neurobiol 69(6):378-391(2009)  
Indraccolo, S., et al. Cancer Res. 69(4):1314-1323(2009)  
Segarra, M., et al. Blood 112(5):1904-1911(2008)

## Images

---



All lanes : Anti-DLL4\_HUMAN at 1:1000 dilution Lane 1: HUVEC whole cell lysate Lane 2: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 75 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

## Citations

---

- [Jagged1 and DLL4 expressions in benign and malignant pancreatic lesions and their clinicopathological significance.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.