

C4BPA Rabbit pAb

Catalog # AP99662

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P04003
Other Accession	P04003
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67033
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C4a
Epitope Specificity	201-300/597
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 8 Sushi (CCP/SCR) domains.
SUBUNIT	Disulfide-linked complex of alpha and beta chains of 3 possible sorts: a 570 kDa complex of 7 alpha chains and 1 beta chain, a 530 kDa homoheptamer of alpha chains or a 500 kDa complex of 6 alpha chains and 1 beta chain. The central body of the alpha chain homopolymer supports tentacles, each with the binding site for C4b at the end.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	C4BPA is a member of a superfamily of proteins composed predominantly of tandemly arrayed short consensus repeats of approximately 60 amino acids. Along with a single, unique beta-chain, seven identical alpha-chains encoded by this gene assemble into the predominant isoform of C4b-binding protein, a multimeric protein that controls activation of the complement cascade through the classical pathway. The genes encoding both alpha and beta chains are located adjacent to each other on human chromosome 1 in the regulator of complement activation gene cluster. Two pseudogenes of this gene are also found in the cluster.

Additional Information

Gene ID	722
Other Names	C4BP; PRP; C4bpa; C4BPA_HUMAN; Proline-rich protein (PRP); C4BPA_MOUSE; complement component 4 binding protein alpha; complement component 4-binding protein, alpha
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	C4BPA
Synonyms	C4BP
Function	Controls the classical pathway of complement activation. It binds as a cofactor to C3b/C4b inactivator (C3bINA), which then hydrolyzes the complement fragment C4b. It also accelerates the degradation of the C4bC2a complex (C3 convertase) by dissociating the complement fragment C2a. Alpha chain binds C4b. It also interacts with anticoagulant protein S and with serum amyloid P component.
Cellular Location	Secreted.
Tissue Location	Chylomicrons in the plasma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.