

HTRA3 Rabbit pAb

Catalog # AP99681

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P83110
Other Accession	P83110
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48608
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HTRA3
Epitope Specificity	61-160/453
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the peptidase S1B family. Contains 1 IGFBP N-terminal domain. Contains 1 Kazal-like domain. Contains 1 PDZ (DHR) domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	HtrA3 is a 453 amino acid secreted protein that contains one PDZ domain, one Kazal-like domain and one IGFBP N-terminal domain. Widely expressed with particularly high expression in ovary, heart, placenta and bladder, HtrA3 functions as a probable serine protease that is thought to play an important role in placental development. HtrA3 is downregulated in ovarian tumors, suggesting an alternative role in tumor suppression. HtrA3 exists as two alternatively spliced isoforms, designated short and long. The gene encoding HtrA3 maps to human chromosome 4, which encodes nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes.

Additional Information

Gene ID	94031
Other Names	Prsp; Tasp; RGD1308120; HTRA3_HUMAN; HTRA3; High-temperature requirement factor A3; Pregnancy-related serine protease; 3.4.21.-; HTRA3_RAT; HtrA serine peptidase 3
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	HTRA3
Synonyms	PRSP
Function	Serine protease that cleaves beta-casein/CSN2 as well as several extracellular matrix (ECM) proteoglycans such as decorin/DCN, biglycan/BGN and fibronectin/FN1. Inhibits signaling mediated by TGF- beta family proteins possibly indirectly by degradation of these ECM proteoglycans (By similarity). May act as a tumor suppressor. Negatively regulates, in vitro, trophoblast invasion during placental development and may be involved in the development of the placenta in vivo. May also have a role in ovarian development, granulosa cell differentiation and luteinization (PubMed: 21321049 , PubMed: 22229724).
Cellular Location	Secreted. Note=Secretion increased during decidualization of endometrial stromal cells
Tissue Location	Widely expressed, with highest levels in both adult and fetal heart, ovary, uterus placenta, and bladder. In the endometrium, expressed in epithelial glands and the stroma. Also present in leukocytes. Isoform 1 is predominant in heart and skeletal muscle, whereas isoform 2 is predominant in placenta and kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.