

CST9 Rabbit pAb

Catalog # AP99753

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q5W186
Other Accession	Q5W186
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18135
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CST9/Cystatin 9
Epitope Specificity	21-120/159
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the cystatin family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The cystatin superfamily is a well-established family of cysteine protease inhibitors. All true cystatins inhibit cysteine peptidases of the papain family, such as cathepsins, while some also inhibit legumain family enzymes. The CRES (cystatin-related epididymal spermatogenic) protein defines a new subgroup in the family 2 cystatins of the cystatin superfamily. CRES proteins lack two of the three consensus sites necessary for the cystatin inhibition of C1 cysteine proteases. Due to their expression patterns, it is likely that CRES proteins may perform unique and tissue-specific functions in the reproductive and neuroendocrine systems. As a member of the CRES subfamily, Cystatin 9, also designated CLM, is a 159 amino acid protein that is expressed in heart, placenta, lung, liver, skeletal muscle and pancreas. Cystatin 9 is thought to play a role in hematopoietic differentiation or inflammation. It has also been shown to be upregulated by LPS in some cancer cell lines, such as promyelocytic leukemia (HL-60) and myelomonocytic leukemia.

Additional Information

Gene ID	128822
Other Names	CLM; CTES7A; CST9_HUMAN; CST9; Cystatin-like molecule; cystatin 9; testatin
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CST9
Function	May be involved in testis development (By similarity). May play a role in hematopoietic differentiation or inflammation (PubMed: 12535658). Has immunomodulatory and antimicrobial functions against Francisella tularensis, a Gram-negative bacteria (PubMed: 23922243).
Cellular Location	Secreted. Note=May be targeted through the Golgi via the secretory pathway
Tissue Location	Expressed in heart, placenta, lung, liver, skeletal muscle and pancreas (PubMed:12535658). Not expressed in brain (PubMed:12535658). Expressed in epididymis, kidney, testis, spinal cord, and thymus with a strong expression in epididymis and kidney and a weak expression in the spinal cord and thymus (PubMed:20565543)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.