

Artemin Rabbit pAb

Catalog # AP99757

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q5T4W7
Other Accession	Q5T4W7
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22878
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Artemin
Epitope Specificity	141-220/220
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the TGF-beta family. GDNF subfamily.
SUBUNIT	Homodimer; disulfide-linked. Binds to RET.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a secreted ligand of the glial cell line-derived neurotrophic factor (GDNF) subfamily and TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This protein signals through the RET receptor and GFR alpha 3 coreceptor, and supports the survival of a number of peripheral neuron populations and at least one population of dopaminergic CNS neurons. This protein has also been shown to promote tumor growth, metastasis, and drug resistance in mammary carcinoma. [provided by RefSeq, Aug 2016]

Additional Information

Gene ID	9048
Other Names	ART; ENOVIN; EVN; NBN; neublastin; ARTN_HUMAN; ARTN; ARTN_MOUSE; ARTN_RAT; artemin; neurotrophic factor
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ARTN {ECO:0000303 PubMed:9883723, ECO:0000312 HGNC:HGNC:727}
Function	Growth factor that supports the survival of sensory and sympathetic peripheral neurons in culture and also supports the survival of dopaminergic neurons of the ventral mid-brain (PubMed: 10583383 , PubMed: 9883723). Acts by binding to its coreceptor, GFRA3, leading to autophosphorylation and activation of the RET receptor (PubMed: 31535977). Strong attractant of gut hematopoietic cells thus promoting the formation Peyer's patch-like structures, a major component of the gut-associated lymphoid tissue (By similarity).
Cellular Location	Secreted.
Tissue Location	Ubiquitous. Expressed at high levels in peripheral tissues including prostate, placenta, pancreas, heart, kidney, pituitary gland, lung and testis. Expressed at low levels in the brain

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.