

WNT10B Rabbit pAb

Catalog # AP99780

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	O00744
Other Accession	O00744
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43000
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human WNT10B
Epitope Specificity	201-300/389
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the Wnt family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It may be involved in breast cancer, and its protein signaling is likely a molecular switch that governs adipogenesis. This protein is 96% identical to the mouse Wnt10b protein at the amino acid level. This gene is clustered with another family member, WNT1, in the chromosome 12q13 region. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	7480
Other Names	SHFM6; STHAG8; WNT-12; Wnt12; WN10B_HUMAN; WNT10B; Protein Wnt-12; WN10B_MOUSE; Wnt-10b; Wnt10; Wnt family member 10B; wingless-type MMTV integration site family, member 10B
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	WNT10B
Synonyms	WNT12
Function	Member of the Wnt ligand gene family that encodes for secreted proteins, which activate the Wnt signaling cascade. Specifically activates canonical Wnt/beta-catenin signaling and thus triggers beta-catenin/LEF/TCF-mediated transcriptional programs. Involved in signaling networks controlling stemness, pluripotency and cell fate decisions. Acts in the immune system, mammary gland, adipose tissue, bone and skin.
Cellular Location	Secreted, extracellular space, extracellular matrix. Secreted
Tissue Location	Detected in most adult tissues. Highest levels were found in heart and skeletal muscle. Low levels are found in brain

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.