

CPN2 Rabbit pAb

Catalog # AP99785

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P22792
Other Accession	P22792
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60557
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CPN2
Epitope Specificity	371-470/545
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 12 LRR (leucine-rich) repeats. Contains 1 LRRCT domain. Contains 1 LRRNT domain.
SUBUNIT	Tetramer of two catalytic chains and two glycosylated inactive chains.
Post-translational modifications	Whether or not any Cys residues participate in intrachain bonds is unknown, but they do not form interchain disulfide bonds with the 50 kDa catalytic subunit.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	CPN2 is a zinc metalloprotease, and cleaves carboxy-terminal arginines and lysines from peptides found in the bloodstream such as complement anaphylatoxins, kinins, and creatine kinase MM (CK-MM). By removing only one amino acid, CPN has the ability to change peptide activity and receptor binding. It is a 280 kDa tetrameric glycoprotein that is synthesized by the liver and secreted into the plasma. It consists of 2 identical 83 kDa regulatory subunits (CPN2) and 2 identical 50 kDa catalytic subunits (CPN1). CPN2, the 83 kDa subunit, binds and stabilizes the catalytic subunit at 37 degrees Celsius and keeps it in circulation. Under some circumstances it may be an allosteric modifier of the catalytic subunit. CPN is a member of a larger family of carboxypeptidases, many of which also cleave arginine and lysine. Because of the highly conserved active sites and the possible redundant functions of carboxypeptidases, it has been difficult to elucidate the role of CPN in disease processes.

Additional Information

Gene ID	1370
Other Names	ACBP; CPN2_HUMAN; CPN2; Carboxypeptidase N 83 kDa chain; Carboxypeptidase N large subunit; Carboxypeptidase N polypeptide 2; Carboxypeptidase N regulatory subunit; carboxypeptidase N subunit 2; carboxypeptidase N, polypeptide 2, 83kD; carboxypeptidase N, polypeptide 2

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CPN2
Synonyms	ACBP
Function	The 83 kDa subunit binds and stabilizes the catalytic subunit at 37 degrees Celsius and keeps it in circulation. Under some circumstances it may be an allosteric modifier of the catalytic subunit.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.