

Stra8 Rabbit pAb

Catalog # AP99902

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q7Z7C7
Other Accession	Q7Z7C7
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36908
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Stra8
Epitope Specificity	201-300/330
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Post-translational modifications	Phosphorylated.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in humans. [provided by RefSeq, Nov 2010]

Additional Information

Gene ID	346673
Other Names	RGD1562852; STRA8_HUMAN; STRA8; STRA8_MOUSE; stimulated by retinoic acid 8; stimulated by retinoic acid gene 8 homolog (mouse); stimulated by retinoic acid 8 homolog (mouse)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	STRA8 (HGNC:30653)
-------------	--------------------------------------

Function	Meiosis-inducer required for the transition into meiosis for both female and male germ cells. In female germ cells, acts downstream of ZGLP1 as a key effector of the meiotic program: required for premeiotic DNA replication and subsequent events in meiotic prophase. During spermatogenesis, next to its role in meiotic initiation, promotes (but is not required for) spermatogonial differentiation. In complex with MEIOSIN, directly activates the transcription of a subset of critical meiotic genes playing a central role in cell-cycle switching from mitosis to meiosis.
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:P70278}. Nucleus {ECO:0000250 UniProtKB:P70278}. Note=Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent {ECO:0000250 UniProtKB:P70278}
Tissue Location	Expressed specifically in testis and fetal ovaries.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.