

EPHX2 Rabbit pAb

Catalog # AP99964

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P34913
Other Accession	P34913
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62616
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human EPX2
Epitope Specificity	351-450/555
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the AB hydrolase superfamily. Epoxide hydrolase family.
SUBUNIT	Homodimer.
Post-translational modifications	The N-terminus is blocked. The covalent modification of cysteine by 15-deoxy-Delta12,14-prostaglandin-J2 is autocatalytic and reversible. It may occur as an alternative to other cysteine modifications, such as S-nitrosylation and S-palmitoylation (Probable).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	EPHX2 belongs to the epoxide hydrolase family. The protein, found in both the cytosol and peroxisomes, binds to specific epoxides and converts them to the corresponding dihydrodiols. Mutations in this gene have been associated with familial hypercholesterolemia. Alternate transcriptional splice variants of this gene have been observed but have not been thoroughly characterized.

Additional Information

Gene ID	2053
Other Names	ABHD20; CEH; SEH; Eph2; sEP; HYES_HUMAN; EPX2; HYES_MOUSE; HYES_RAT; epoxide hydrolase 2; epoxide hydrolase 2, cytoplasmic; soluble epoxide hydrolase; bifunctional epoxide hydrolase
Dilution	WB=1:500-2000, IHC-P=1:100-500, IHC-F=1:100-500, IF=1:100-500, ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	EPHX2 (HGNC:3402)
Function	Bifunctional enzyme (PubMed: 12574510). The C-terminal domain has epoxide hydrolase activity and acts on epoxides (alkene oxides, oxiranes) and arene oxides (PubMed: 12574510 , PubMed: 12869654 , PubMed: 22798687). Plays a role in xenobiotic metabolism by degrading potentially toxic epoxides (By similarity). Also determines steady- state levels of physiological mediators (PubMed: 12574510 , PubMed: 12869654 , PubMed: 21217101 , PubMed: 22798687).
Cellular Location	Cytoplasm. Peroxisome.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.