

# Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (conatumumab)

**Recombinant Antibody** Catalog # APR10023

### **Product Information**

**Application** FC, Kinetics, Animal Model

**Primary Accession** 014763

Reactivity Human, Mouse Clonality Monoclonal Isotype IgG1

**Calculated MW** 47878

#### **Additional Information**

Target/Specificity TNFRSF10B / TRAILR2 / CD262

**Endotoxin** 

Conjugation Unconjugated

**Expression system** CHO Cell

**Format** Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

#### **Protein Information**

TNFRSF10B Name

**Synonyms** DR5, KILLER, TRAILR2, TRICK2, ZTNFR9

**Function** Receptor for the cytotoxic ligand TNFSF10/TRAIL (PubMed: 10549288). The

> adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the

activation of NF-kappa-B. Essential for ER stress-induced apoptosis.


**Cellular Location** Membrane; Single-pass type I membrane protein.

**Tissue Location** Widely expressed in adult and fetal tissues; very highly expressed in tumor


> cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and

throughout the intestinal tract; not detectable in brain


## **Images**



Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (conatumumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (conatumumab)is more than 98.9%, determined by SEC-HPLC.



Immobilized Cyno DR5 His at 2 μg/mL can bind Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (conatumumab),EC50=0.1119 μg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.