

# Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (drozitumab)

Recombinant Antibody  
Catalog # APR10025

## Product Information

---

<b>Application</b>	FC, Kinetics, Animal Model
<b>Primary Accession</b>	<a href="#">O14763</a>
<b>Reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	47878

## Additional Information

---


<b>Target/Specificity</b>	TNFRSF10B / TRAILR2 / CD262
<b>Endotoxin Conjugation</b>	Unconjugated
<b>Expression system</b>	CHO Cell
<b>Format</b>	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

## Protein Information


---

<b>Name</b>	TNFRSF10B
<b>Synonyms</b>	DR5, KILLER, TRAILR2, TRICK2, ZTNFR9
<b>Function</b>	Receptor for the cytotoxic ligand TNFSF10/TRAIL (PubMed: <a href="#">10549288</a> ). The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.
<b>Cellular Location</b>	Membrane; Single-pass type I membrane protein.
<b>Tissue Location</b>	Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and throughout the intestinal tract; not detectable in brain


# Images



Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (drozitumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (drozitumab) is more than 99.68% ,determined by SEC-HPLC.



Immobilized Cynomolgus TRAIL R2/DR5/ TNFRSF10B Protein, His Tag at 2 µg/mL can bind Anti-TNFRSF10B / TRAILR2 / CD262 Reference Antibody (drozitumab), EC50=0.1257 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.