

Anti-TNFRSF5 / CD40 Reference Antibody (bleselumab)

Recombinant Antibody

Catalog # APR10028

Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	P25942
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG4PE
Calculated MW	30619

Additional Information



Target/Specificity	TNFRSF5 / CD40
Endotoxin	
Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Protein Information


Name	CD40
Synonyms	TNFRSF5
Function	Receptor for TNFSF5/CD40LG (PubMed: 31331973). Transduces TRAF6- and MAP3K8-mediated signals that activate ERK in macrophages and B cells, leading to induction of immunoglobulin secretion (By similarity).
Cellular Location	[Isoform I]: Cell membrane; Single-pass type I membrane protein
Tissue Location	B-cells and in primary carcinomas.

Images


Anti-TNFRSF5 / CD40 Reference Antibody (bleselumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%




The purity of Anti-TNFRSF5 / CD40 Reference Antibody (bleselumab) is more than 99.21% ,determined by SEC-HPLC.



Immobilized human TNFRSF5 / CD40 His at 2 µg/mL can bind Anti-TNFRSF5 / CD40 Reference Antibody (bleselumab), EC₅₀=0.0252 µg/mL.



Human CD40 CHO cells were stained with Anti-TNFRSF5 / CD40 Reference Antibody (bleselumab) and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC₇₇=0.6311 µg/mL



Anti-TNFRSF5 / CD40 Reference Antibody (bleselumab) FACS Blocking was evaluated using CCRF-CEM. The IC₅₀ was approximately 1.359 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.