

Anti-TNFRSF7 / CD27 Reference Antibody (varlilumab)

Recombinant Antibody Catalog # APR10089

Product Information

Application FC, Kinetics, Animal Model

Primary Accession
Reactivity
Human
Clonality
Monoclonal
Isotype
IgG1
Calculated MW
29137

Additional Information

Target/Specificity TNFRSF7 / CD27

Endotoxin

Conjugation Unconjugated

Expression system CHO Cell

Format Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

Protein Information

Name CD27 (<u>HGNC:11922</u>)

Function Costimulatory immune-checkpoint receptor expressed at the surface of


T-cells, NK-cells and B-cells which binds to and is activated by its ligand CD70/CD27L expressed by B-cells (PubMed:28011863). The CD70-CD27 signaling pathway mediates antigen- specific T-cell activation and expansion which in turn provides immune surveillance of B-cells (PubMed:28011863). Mechanistically, CD70 ligation activates the TRAF2-PTPN6 axis that subsequently inhibits LCK phosphorylation to promote phenotypic and transcriptional adaptations of T-cell memory (PubMed:38354704). In addition, activation by CD70 on early progenitor cells provides a negative feedback signal to leukocyte differentiation during immune activation and thus modulates hematopoiesis (By similarity). Negatively regulates the function of

Th2 lymphocytes in the adipose tissue (By similarity).


Cellular Location Cell membrane; Single-pass type I membrane protein

Tissue Location Found in most T-lymphocytes.


Images




Anti-TNFRSF7 / CD27 Reference Antibody (varlilumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-TNFRSF7 / CD27 Reference Antibody (varlilumab)is more than 100% ,determined by SEC-HPLC.



Immobilized human CD27 FC at 2 µg/mL can bind Anti-TNFRSF7 / CD27 Reference Antibody (varlilumab),EC50=0.01912 µg/mL



HumanCD27 HEK293 cells were stained with Anti-TNFRSF7 / CD27 Reference Antibody (varlilumab) and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC142=0.095 ug/mL

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.