10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-FOLR1 / FRA Reference Antibody (mirvetuximab)

Recombinant Antibody
Catalog # APR10114

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P15328
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	29819

Additional Information


Target/Specificity	FOLR1 / FRA
Endotoxin	
Conjugation	Unconjugated
Expression system	CHO
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Protein Information


Name	FOLR1
Synonyms	FOLR
Function	Binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells (PubMed:19074442, PubMed:23851396, PubMed:23934049, PubMed:2527252, PubMed:8033114, PubMed:8567728). Has high affinity for folate and folic acid analogs at neutral pH (PubMed:23851396, PubMed:23934049, PubMed:2527252, PubMed:8033114, PubMed:8567728). Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release (PubMed:8567728). Required for normal embryonic development and normal cell proliferation (By similarity).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor Apical cell membrane; Lipid-anchor, GPI- anchor Basolateral cell membrane; Lipid-anchor, GPI-like-anchor. Secreted Cytoplasmic vesicle. Cytoplasmic vesicle, clathrin-coated vesicle. Endosome. Note=Endocytosed into cytoplasmic vesicles and then recycled to the cell membrane

Tissue Location	Primarily expressed in tissues of epithelial origin. Expression is increased in malignant tissues. Expressed in kidney, lung and cerebellum. Detected in placenta and thymus epithelium.


Images




Anti-FOLR1 / FRA Reference Antibody (mirvetuximab) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.




The purity of Anti-FOLR1 / FRA Reference Antibody (mirvetuximab) is 100%, determined by SEC-HPLC.



Immobilized human FRa His at 2 ug/mL can bind Anti-FOLR1 / FRA Reference Antibody (mirvetuximab), EC50=0.01123 ug/mL.



SKOV3 cells were stained with Anti-FOLR1 / FRA Reference Antibody (mirvetuximab) and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC50=0.03238 ug/mL.



The endocytosis ratio mirvetuximab by Human Frα HEK 293 increased with the increase of antibody concentration, and the Internalization Rate (%) 75% at antibody concentration of 55 ng/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.