

Anti-ICOS / CD278 Reference Antibody (feladilimab)

Recombinant Antibody

Catalog # APR10123

Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	Q9Y6W8
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG4PE
Calculated MW	22625

Additional Information


Target/Specificity	ICOS / CD278
Endotoxin	
Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Protein Information


Name	ICOS
Synonyms	AILIM
Function	Stimulatory receptor expressed in activated or antigen-experienced T-cells that plays an important role in the immune response (PubMed: 9930702). Upon binding to its ligand ICOSL expressed on antigen presenting cells (APCs), delivers costimulatory signals that enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines including IL10, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells (PubMed: 33033255). Also acts as a costimulatory receptor critical for the differentiation of T follicular regulatory cells upon immune challenges such as viral infection (PubMed: 27135603). Mechanistically, potentiates TCR-induced calcium flux by augmenting PLCG1 activation and actin remodeling (By similarity). In addition, activates PI3K signaling pathways independently of calcium flux (PubMed: 30523347). Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40-mediated class switching of immunoglobulin isotypes (By similarity).

Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus


Images



Anti-ICOS / CD278 Reference Antibody (feladilimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-ICOS / CD278 Reference Antibody (feladilimab)is more than 99.12% ,determined by SEC-HPLC.



Immobilized human ICOS FC at 2 µg/mL can bind Anti-ICOS / CD278 Reference Antibody (feladilimab),EC50=0.09127 µg/mL

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.