

# Anti-CXCL8 / IL-8 Reference Antibody (HuMax-IL8)

Recombinant Antibody  
Catalog # APR10223

## Product Information

---

<b>Application</b>	FC, Kinetics, Animal Model
<b>Primary Accession</b>	<a href="#">P10145</a>
<b>Reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	11098

## Additional Information


---

<b>Target/Specificity</b>	CXCL8 / IL-8
<b>Endotoxin Conjugation</b>	Unconjugated
<b>Expression system</b>	CHO Cell
<b>Format</b>	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.


## Protein Information

---


<b>Name</b>	CXCL8
<b>Synonyms</b>	IL8
<b>Function</b>	Chemotactic factor that mediates inflammatory response by attracting neutrophils, basophils, and T-cells to clear pathogens and protect the host from infection (PubMed: <a href="#">18692776</a> , PubMed: <a href="#">7636208</a> ). Also plays an important role in neutrophil activation (PubMed: <a href="#">2145175</a> , PubMed: <a href="#">9623510</a> ). Released in response to an inflammatory stimulus, exerts its effect by binding to the G-protein-coupled receptors CXCR1 and CXCR2, primarily found in neutrophils, monocytes and endothelial cells (PubMed: <a href="#">1840701</a> , PubMed: <a href="#">1891716</a> ). G-protein heterotrimer (alpha, beta, gamma subunits) constitutively binds to CXCR1/CXCR2 receptor and activation by IL8 leads to beta and gamma subunits release from Galpha (GNAI2 in neutrophils) and activation of several downstream signaling pathways including PI3K and MAPK pathways (PubMed: <a href="#">11971003</a> , PubMed: <a href="#">8662698</a> ).
<b>Cellular Location</b>	Secreted.



Anti-CXCL8 / IL-8 Reference Antibody (HuMax-IL8) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 91.8%



The purity of Anti-CXCL8 / IL-8 Reference Antibody (HuMax-IL8) is more than 91.32% ,determined by SEC-HPLC.



Immobilized human IL 8 His at 2 µg/mL can bind Anti-CXCL8 / IL-8 Reference Antibody (HuMax-IL8),EC50=0.0427 µg/mL

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.