

Anti-NKG2D / CD314 Reference Antibody (tesnatilimab)

Recombinant Antibody

Catalog # APR10241

Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	P26718
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG4
Calculated MW	25304

Additional Information

Target/Specificity	NKG2D / CD314
Endotoxin	
Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.


Protein Information

Name	KLrk1
Synonyms	D12S2489E, NKG2D
Function	Functions as an activating and costimulatory receptor involved in immunosurveillance upon binding to various cellular stress- inducible ligands displayed at the surface of autologous tumor cells and virus-infected cells. Provides both stimulatory and costimulatory innate immune responses on activated killer (NK) cells, leading to cytotoxic activity. Acts as a costimulatory receptor for T-cell receptor (TCR) in CD8(+) T-cell-mediated adaptive immune responses by amplifying T-cell activation. Stimulates perforin-mediated elimination of ligand-expressing tumor cells. Signaling involves calcium influx, culminating in the expression of TNF-alpha. Participates in NK cell- mediated bone marrow graft rejection. May play a regulatory role in differentiation and survival of NK cells. Binds to ligands belonging to various subfamilies of MHC class I-related glycoproteins including MICA, MICB, RAET1E, RAET1G, RAET1L/ULBP6, ULBP1, ULBP2, ULBP3 (ULBP2>ULBP1>ULBP3) and ULBP4.
Cellular Location	Cell membrane; Single-pass type II membrane protein Note=Colocalized with HCST on the cell surface


Tissue Location

Expressed in natural killer (NK) cells, CD8(+) alpha-beta and gamma-delta T-cells. Expressed on essentially all CD56+CD3- NK cells from freshly isolated PBMC. Expressed in interferon- producing killer dendritic cells (IKDCs).


Images



Anti-NKG2D / CD314 Reference Antibody (tesnatilimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-NKG2D / CD314 Reference Antibody (tesnatilimab) is more than 100% ,determined by SEC-HPLC.



Immobilized human NKG2D His at 2 µg/mL can bind Anti-NKG2D / CD314 Reference Antibody (tesnatilimab),EC50=0.0209 µg/mL

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.