

Anti-CCR2 / CD192 Reference Antibody (plozalizumab)

Recombinant Antibody

Catalog # APR10318

Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	P41597
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	41915

Additional Information

Target/Specificity	CCR2 / CD192
Endotoxin	
Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Protein Information

Name	CCR2
Synonyms	CMKBR2
Function	<p>Key functional receptor for CCL2 but can also bind CCL7, and CCL12 (PubMed:23408426, PubMed:38157855, PubMed:8048929, PubMed:8146186). Also transduces signaling mediated by CCL13 (PubMed:38157855). Its binding with CCL2 on monocytes and macrophages mediates chemotaxis and migration induction through the activation of the PI3K cascade, the small G protein Rac and lamellipodium protrusion (PubMed:38157855). Also acts as a receptor for the beta-defensin DEFB106A/DEFB106B (PubMed:23938203). Regulates the expression of T-cell inflammatory cytokines and T-cell differentiation, promoting the differentiation of T-cells into T-helper 17 cells (Th17) during inflammation (By similarity). Facilitates the export of mature thymocytes by enhancing directional movement of thymocytes to sphingosine-1-phosphate stimulation and up-regulation of S1P1R expression; signals through the JAK-STAT pathway to regulate FOXO1 activity leading to an increased expression of S1P1R (By similarity). Plays an important role in mediating peripheral nerve injury-induced neuropathic pain (By similarity). Increases NMDA-mediated synaptic transmission in both dopamine D1 and D2 receptor-containing neurons, which may be caused by</p>

MAPK/ERK-dependent phosphorylation of GRIN2B/NMDAR2B (By similarity). Mediates the recruitment of macrophages and monocytes to the injury site following brain injury (By similarity).


Cellular Location

Cell membrane; Multi-pass membrane protein. Note=The chemoattractant receptors are distributed throughout the cell surface; after stimulation with a ligand, such as CCL2, they are rapidly recruited into microdomain clusters at the cell membrane.


Tissue Location

Expressed by monocytes and IL2-activated NK cells (PubMed:9058802). Abundantly expressed on CD14+/CD16- monocytes and weakly on CD14+/CD16+ monocytes, type 2 dendritic cells (DCs) and plasmacytoid DCs (at protein level) (PubMed:38157855)

Images



Anti-CCR2 / CD192 Reference Antibody (plozalizumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-CCR2 / CD192 Reference Antibody (plozalizumab) is more than 99.45% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.