

Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab)

Recombinant Antibody Catalog # APR10343

Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	<u>P29965</u>
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	29274

Additional Information

Target/Specificity	TNFSF5 / CD40L / CD154
Endotoxin Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Protein Information

Name Synonyms	CD40LG CD40L, TNFSF5, TRAP
Function	Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed: <u>1280226</u> , PubMed: <u>31331973</u>). Costimulates T-cell proliferation and cytokine production (PubMed: <u>8617933</u>). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed: <u>8617933</u>). Induces the activation of NF-kappa-B (PubMed: <u>15067037</u> , PubMed: <u>31331973</u>). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed: <u>15067037</u>). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed: <u>15067037</u>). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).
Cellular Location	Cell membrane; Single-pass type II membrane protein. Cell surface
Tissue Location	Specifically expressed on activated CD4+ T- lymphocytes

Images



Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%

The purity of Anti-TNFSF5 / CD40L / CD154 Reference Antibody (ruplizumab)is more than 99% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.