

# Anti-MUC18 / MCAM / CD146 Reference Antibody (imaprelimab)

Recombinant Antibody Catalog # APR10374

### **Product Information**

**Application** FC, Kinetics, Animal Model

Primary Accession P43121

Reactivity Human, Mouse
Clonality Monoclonal
Isotype IgG1

Calculated MW 71607

#### **Additional Information**

Target/Specificity MUC18 / MCAM / CD146

**Endotoxin** 

**Conjugation** Unconjugated

**Expression system** CHO Cell

Format Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

#### **Protein Information**

Name MCAM

Synonyms MUC18

**Function** Plays a role in cell adhesion, and in cohesion of the endothelial monolayer at

intercellular junctions in vascular tissue. Its expression may allow melanoma cells to interact with cellular elements of the vascular system, thereby enhancing hematogeneous tumor spread. Could be an adhesion molecule active in neural crest cells during embryonic development. Acts as a surface receptor that triggers tyrosine phosphorylation of FYN and PTK2/FAK1, and a

transient increase in the intracellular calcium concentration.


**Cellular Location** Membrane; Single-pass type I membrane protein.

**Tissue Location** Detected in endothelial cells in vascular tissue throughout the body. May


appear at the surface of neural crest cells during their embryonic migration. Appears to be limited to vascular smooth muscle in normal adult tissues. Associated with tumor progression and the development of metastasis in human malignant melanoma. Expressed most strongly on metastatic lesions

and advanced primary tumors and is only rarely detected in benign melanocytic nevi and thin primary melanomas with a low probability of metastasis

## **Images**



Anti-MUC18 / MCAM / CD146 Reference Antibody (imaprelimab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-MUC18 / MCAM / CD146 Reference Antibody (imaprelimab)is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.