10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-ERBB3 / HER3 Reference Antibody (seribantumab)

Recombinant Antibody
Catalog # APR10434

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P21860
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG2SA
Calculated MW	148098

Additional Information


Target/Specificity	ERBB3 / HER3
Endotoxin	
Conjugation	Unconjugated
Expression system	CHO
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Protein Information


Name	ERBB3
Synonyms	HER3
Function	Tyrosine kinase receptor that, despite substitutions of key catalytic residues in the kinase domain, retains residual catalytic activity enabling autophosphorylation (PubMed:9148746, PubMed:20351256). Functions primarily as a heterodimer, preferentially partnering with ERBB2/HER2, which phosphorylates ERBB3 on specific tyrosine residues and provides the kinase activity required for downstream signaling (PubMed:22983903). Binds neuregulins NRG1 and NRG2 through its extracellular domain, which induces conformational change from a tethered autoinhibited state to an extended conformation competent for dimerization (PubMed:20682778). Upon heterodimerization and trans-phosphorylation, acts as the most efficient activator of the PI3K/AKT pathway within the ErbB family (PubMed:22983903). Also activates the RAS/MAPK/ERK cascade to promote cell proliferation (PubMed:22983903). Plays an essential role in cardiac development; required for normal formation of the endocardial cushion and ventricular trabeculation during embryogenesis (Potential). Critical for peripheral nervous system development, where neuregulin-1/ERBB3 signaling drives Schwann cell survival, proliferation, and differentiation, and is required for

	axonal myelination (By similarity). Involved in epithelial morphogenesis and mammary gland development (By similarity). Also involved in the regulation of myeloid cell differentiation (PubMed:27416908). Participates in a regulatory feedback loop whereby pathway activation induces transcriptional up-regulation of ERBB3 expression (PubMed:21368164). Participates in vesicular sorting by stabilizing the ARF6-GGA3-RABEP1 endosomal sorting complex, to promote early recycling of TFRC and ITGB1 (PubMed:41348103). May also be activated by additional ligands including CSPG5 and EGFL6; the latter triggers assembly of an ERBB2/ERBB3/ITGB3 signaling complex leading to ERK phosphorylation and cell proliferation (PubMed:15358134, PubMed:41151154).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Epithelial tissues and brain.

Images



Anti-ERBB3 / HER3 Reference Antibody (seribantumab) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 90%.



The purity of Anti-ERBB3 / HER3 Reference Antibody (seribantumab) is more than 95%, determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.