10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-FOLR1 / FRA Reference Antibody (Mirvetuximab soravtansine)

Recombinant Antibody
Catalog # APR10452

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P15328
Reactivity	Human
Clonality	ADC
Isotype	IgG1
Calculated MW	29819

Additional Information


Target/Specificity	FOLR1 / FRA
Endotoxin Conjugation	DM4
Expression system	CHO
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

Protein Information


Name	FOLR1
Synonyms	FOLR
Function	Binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells (PubMed:19074442, PubMed:23851396, PubMed:23934049, PubMed:2527252, PubMed:8033114, PubMed:8567728). Has high affinity for folate and folic acid analogs at neutral pH (PubMed:23851396, PubMed:23934049, PubMed:2527252, PubMed:8033114, PubMed:8567728). Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release (PubMed:8567728). Required for normal embryonic development and normal cell proliferation (By similarity).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor Apical cell membrane; Lipid-anchor, GPI- anchor Basolateral cell membrane; Lipid-anchor, GPI-like-anchor. Secreted Cytoplasmic vesicle. Cytoplasmic vesicle, clathrin-coated vesicle. Endosome. Note=Endocytosed into cytoplasmic vesicles and then recycled to

	the cell membrane
Tissue Location	Primarily expressed in tissues of epithelial origin. Expression is increased in malignant tissues. Expressed in kidney, lung and cerebellum. Detected in placenta and thymus epithelium.


Images



Anti-FOLR1 / FRA Reference Antibody (Mirvetuximab soravtansine) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.



The purity of Anti-FOLR1 / FRA Reference Antibody (Mirvetuximab soravtansine) is more than 95%, determined by SEC-HPLC.



Immobilized human FRA, His Tag at 2 ug/mL can bind Anti-FOLR1 / FRA Reference Antibody (Mirvetuximab soravtansine), EC50 is 0.009438 ug/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.