10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-IL-13 Reference Antibody (IMA-026)

Recombinant Antibody
Catalog # APR10511

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P35225
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	15788

Additional Information

Target/Specificity	IL-13
Endotoxin	
Conjugation	Unconjugated
Expression system	CHO
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.


Protein Information

Name	IL13
Synonyms	NC30
Function	Cytokine that plays important roles in allergic inflammation and immune response to parasite infection (PubMed:8096327, PubMed:8097324). Synergizes with IL2 in regulating interferon-gamma synthesis (PubMed:8096327). Stimulates B-cell proliferation, and activation of eosinophils, basophils, and mast cells (PubMed:7903680, PubMed:8759755). Plays an important role in controlling IL33 activity by modulating the production of transmembrane and soluble forms of interleukin-1 receptor-like 1/IL1RL1 (By similarity). Displays the capacity to antagonize Th1-driven proinflammatory immune response and down-regulates synthesis of many proinflammatory cytokines including IL1, IL6, IL10, IL12 and TNF through a mechanism that partially involves suppression of NF-kappa-B (By similarity). Also functions on nonhematopoietic cells, including endothelial cells where it induces vascular cell adhesion protein 1/VCAM1, which is important in the recruitment of eosinophils (PubMed:8639787). Exerts its biological effects through its receptors which comprises the IL4R chain and the IL13RA1 chain, to activate JAK1 and TYK2, leading to the activation of STAT6 (PubMed:9013879). Aside from IL13RA1, another receptor IL13RA2 acts


as a high affinity decoy for IL13 and mediates internalization and depletion of extracellular IL13 (PubMed:21622864). Induces inflammatory response in esophageal epithelium which results in a dysregulation of desmosome intercellular adhesion junctions, via down-regulation of DSG1 (PubMed:24220297).

Cellular Location Secreted.

Images



Anti-IL-13 Reference Antibody (IMA-026) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.



The purity of Anti-IL-13 Reference Antibody (IMA-026) is more than 95%, determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.