

# Anti-ACVR1 / ALK-2 Reference Antibody (DS-6016a)

Recombinant Antibody Catalog # APR10588

# **Product Information**

Application	FC, Kinetics, Animal Model
Primary Accession	<u>Q04771</u>
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	57153

### **Additional Information**

Target/Specificity	ACVR1 / ALK-2
Endotoxin Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.

# **Protein Information**

Name	ACVR1
Synonyms	ACVRLK2
Function	Bone morphogenetic protein (BMP) type I receptor that is involved in a wide variety of biological processes, including bone, heart, cartilage, nervous, and reproductive system development and regulation (PubMed: <u>20628059</u> , PubMed: <u>22977237</u> ). As a type I receptor, forms heterotetrameric receptor complexes with the type II receptors AMHR2, ACVR2A or ACVR2B (PubMed: <u>17911401</u> ). Upon binding of ligands such as BMP7 or GDF2/BMP9 to the heteromeric complexes, type II receptors transphosphorylate ACVR1 intracellular domain (PubMed: <u>25354296</u> ). In turn, ACVR1 kinase domain is activated and subsequently phosphorylates SMAD1/5/8 proteins that transduce the signal (PubMed: <u>9748228</u> ). In addition to its role in mediating BMP pathway-specific signaling, suppresses TGFbeta/activin pathway signaling by interfering with the binding of activin to its type II receptor (PubMed: <u>17911401</u> ). Besides canonical SMAD signaling, can activate non-canonical pathways such as p38 mitogen-activated protein kinases/MAPKs (By similarity). May promote the expression of HAMP, potentially via its interaction with BMP6 (By similarity).

#### **Cellular Location**

Membrane; Single-pass type I membrane protein.

**Tissue Location** 

Expressed in normal parenchymal cells, endothelial cells, fibroblasts and tumor-derived epithelial cells

#### Images



Anti-ACVR1 / ALK-2 Reference Antibody (DS-6016a) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%

The purity of Anti-ACVR1 / ALK-2 Reference Antibody (DS-6016a)is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.