



# Anti-TNFSF2 / TNFa Reference Antibody (certolizumAb)

Recombinant Antibody Catalog # APR10617

#### **Product Information**

**Application** FC, Kinetics, Animal Model

Primary Accession P01375

Reactivity Human, Mouse
Clonality Monoclonal
Isotype IgG1

Isotype IgG1
Calculated MW 25644

### **Additional Information**

Target/Specificity TNFSF2 / TNFa

**Endotoxin** 

**Conjugation** Unconjugated

**Expression system** CHO Cell

**Format** Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

### **Protein Information**

Name TNF

Synonyms TNFA, TNFSF2

**Function** Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly

secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under

certain conditions it can stimulate cell proliferation and induce cell

differentiation. Impairs regulatory T- cells (Treg) function in individuals with rheumatoid arthritis via FOXP3 dephosphorylation. Up-regulates the

expression of protein phosphatase 1 (PP1), which dephosphorylates the key 'Ser-418' residue of FOXP3, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed: 23396208). Key mediator of cell death in the anticancer action of BCG-stimulated neutrophils in combination with

DIABLO/SMAC mimetic in the RT4v6 bladder cancer cell line


(PubMed:<u>16829952</u>, PubMed:<u>22517918</u>, PubMed:<u>23396208</u>). Induces insulin resistance in adipocytes via inhibition of insulin-induced IRS1 tyrosine phosphorylation and insulin-induced glucose uptake. Induces GKAP42 protein degradation in adipocytes which is partially responsible for TNF-induced insulin resistance (By similarity). Plays a role in angiogenesis by inducing VEGF

production synergistically with IL1B and IL6 (PubMed: 12794819). Promotes osteoclastogenesis and therefore mediates bone resorption (By similarity).


#### **Cellular Location**

Cell membrane; Single-pass type II membrane protein [Tumor necrosis factor, soluble form]: Secreted [C-domain 2]: Secreted.

## **Images**



Anti-TNFSF2 / TNFa Reference Antibody (certolizumAb) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-TNFSF2 / TNFa Reference Antibody (certolizumAb)is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.