

Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151)

Recombinant Antibody
Catalog # APR10799

Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	P48509
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	28295

Additional Information

Target/Specificity	CD151
Endotoxin Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Protein Information

Name	CD151
Synonyms	TSPAN24
Function	Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling. Plays a role in various cellular and molecular mechanism through its association with both integrin and non-integrin proteins. These interactions facilitate critical cellular functions, including cell-to-cell communication, wound healing, platelet aggregation, trafficking, cell motility, and angiogenesis (PubMed: 17045834 , PubMed: 24723389 , PubMed: 31488507). Via interaction with JAM-A/F11R and integrin ITGA3:ITGB1, promotes the recruitment of signaling molecules such as RAC1, CDC42 and RhoGTPases to facilitate the polarization of epithelial cells and the reorganization of the actin cytoskeleton, which are critical steps in cell migration process (PubMed: 22843693 , PubMed: 35067832). Regulates the glycosylation pattern of ITGA3:ITGB1 thereby modulating its activity (PubMed: 18852263). Plays an essential role in the maintenance of central laminin-binding integrin ITGA6:ITGB4-containing adhesion complexes

(PubMed:[31488507](#)). Essential for the proper assembly of the glomerular and tubular basement membranes in kidney (PubMed:[15265795](#)). Contributes to T-cell activation by modulating integrin signaling leading to activation of downstream targets PTK2 and MAPK1/MAPK3 (PubMed:[24723389](#)).


Cellular Location

Cell membrane; Multi-pass membrane protein Note=Relocalizes to the immune synapse in T-cells upon activation


Tissue Location

Expressed in a variety of tissues including vascular endothelium and epidermis. Expressed on erythroid cells, with a higher level of expression in erythroid precursors than on mature erythrocytes (PubMed:15265795). Acts as a sensitive T-cell activation marker (PubMed:32978478).

Images



Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-CD151 Reference Antibody (Pierre Fabre patent anti-CD151) is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.