

Anti-CXCL4 / PF4 Reference Antibody (U.Penn. patent anti-PF4)

Recombinant Antibody
Catalog # APR10872

Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	P02776
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	10845

Additional Information

Target/Specificity	CXCL4 / PF4
Endotoxin Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Protein Information


Name	PF4
Synonyms	CXCL4, SCYB4
Function	Chemokine released during platelet aggregation that plays a role in different biological processes including hematopoiesis, cell proliferation, differentiation, and activation (PubMed: 29930254 , PubMed: 9531587). Acts via different functional receptors including CCR1, CXCR3A or CXCR3B (PubMed: 18174362 , PubMed: 29930254). Upon interaction with CXCR3A receptor, induces activated T-lymphocytes migration mediated via downstream Ras/extracellular signal-regulated kinase (ERK) signaling (PubMed: 18174362 , PubMed: 24469069). Neutralizes the anticoagulant effect of heparin by binding more strongly to heparin than to the chondroitin-4-sulfate chains of the carrier molecule. Plays a role in the inhibition of hematopoiesis and in the maintenance of hematopoietic stem cell (HSC) quiescence (PubMed: 9531587). Chemotactic for neutrophils and monocytes via CCR1 (PubMed: 29930254). Inhibits endothelial cell proliferation. In cooperation with toll-like receptor 8/TLR8, induces chromatin remodeling and activates inflammatory gene expression via the TBK1-IRF5

axis (PubMed:[35701499](#)). In addition, induces myofibroblast differentiation and collagen synthesis in different precursor cells, including endothelial cells, by stimulating endothelial-to-mesenchymal transition (PubMed:[34986347](#)). Interacts with thrombomodulin/THBD to enhance the activation of protein C and thus potentiates its anticoagulant activity (PubMed:[9395524](#)).


Cellular Location

Secreted.

Images



Anti-CXCL4 / PF4 Reference Antibody (U.Penn. patent anti-PF4) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-CXCL4 / PF4 Reference Antibody (U.Penn. patent anti-PF4) is more than 95%, determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.