



Anti-CXCL8 / IL-8 Reference Antibody (Genentech patent anti-IL-8)

Recombinant Antibody Catalog # APR10873

Product Information

FC, Kinetics, Animal Model **Application**

Primary Accession P10145 Reactivity Human Clonality Monoclonal Isotype IgG1 **Calculated MW** 11098

Additional Information

Target/Specificity CXCL8 / IL-8

Endotoxin

Conjugation Unconjugated

Expression system CHO Cell

Format Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

Protein Information

Name CXCL8

Synonyms IL8

Function Chemotactic factor that mediates inflammatory response by attracting

neutrophils, basophils, and T-cells to clear pathogens and protect the host from infection (PubMed: 18692776, PubMed: 7636208). Also plays an

important role in neutrophil activation (PubMed:2145175, PubMed:9623510). Released in response to an inflammatory stimulus, exerts its effect by binding to the G-protein-coupled receptors CXCR1 and CXCR2, primarily found in


neutrophils, monocytes and endothelial cells (PubMed: 1840701,

PubMed: 1891716). G-protein heterotrimer (alpha, beta, gamma subunits) constitutively binds to CXCR1/CXCR2 receptor and activation by IL8 leads to beta and gamma subunits release from Galpha (GNAI2 in neutrophils) and activation of several downstream signaling pathways including PI3K and


MAPK pathways (PubMed: 11971003, PubMed: 8662698).

Cellular Location Secreted.

Images



Anti-CXCL8 / IL-8 Reference Antibody (Genentech patent anti-IL-8) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-CXCL8 / IL-8 Reference Antibody (Genentech patent anti-IL-8)is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.