

# Anti-Fibronectin Reference Antibody (L19-TNF)

Recombinant Antibody Catalog # APR10910

### **Product Information**

**Application** FC, Kinetics, Animal Model

Primary Accession
Reactivity
Human
Clonality
Monoclonal
Isotype
IgG4
Calculated MW
272320

#### **Additional Information**

Target/Specificity Fibronectin

**Endotoxin** 

**Conjugation** Unconjugated

**Expression system** CHO Cell

**Format** Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

## **Protein Information**

Name FN1 ( HGNC:3778)

Synonyms FN

**Function** Fibronectins bind cell surfaces and various compounds including collagen,

fibrin, heparin, DNA, and actin (PubMed:3024962, PubMed:3593230, PubMed:3900070, PubMed:7989369). Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell

shape (PubMed:3024962, PubMed:3593230, PubMed:3900070,

PubMed: 7989369). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). Participates in the regulation of type I collagen deposition by osteoblasts (By similarity). Acts as a ligand for the LILRB4

receptor, inhibiting FCGR1A/CD64-mediated monocyte activation

(PubMed:34089617).

**Cellular Location** Secreted, extracellular space, extracellular matrix. Secreted


{ECO:0000250 | UniProtKB:P11276}

**Tissue Location** Expressed in the inner limiting membrane and around blood vessels in the


retina (at protein level) (PubMed:29777959) Plasma FN (soluble dimeric form)

is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine (PubMed:17614963).

# **Images**



Anti-Fibronectin Reference Antibody (L19-TNF) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-Fibronectin Reference Antibody (L19-TNF)is more than 98.36% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.