10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



Anti-FOLR1 / FRA Reference Antibody (Dompe patent anti-FOLR1)

Recombinant Antibody Catalog # APR10913

Product Information

Application FC, Kinetics, Animal Model

29819

Primary Accession
Reactivity
Human
Clonality
Monoclonal
Isotype
IgG1

Additional Information

Target/Specificity FOLR1 / FRA

Endotoxin

Calculated MW

Conjugation Unconjugated

Expression system CHO Cell

Format Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

Protein Information

Name FOLR1

Synonyms FOLR

Function Binds to folate and reduced folic acid derivatives and mediates delivery of

5-methyltetrahydrofolate and folate analogs into the interior of cells

(PubMed:19074442, PubMed:23851396, PubMed:23934049,

PubMed: 2527252, PubMed: 8033114, PubMed: 8567728). Has high affinity for

folate and folic acid analogs at neutral pH (PubMed: 23851396,

PubMed:23934049, PubMed:2527252, PubMed:8033114, PubMed:8567728).

Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release (PubMed:8567728). Required for normal embryonic

development and normal cell proliferation (By similarity).

Cellular Location Cell membrane; Lipid-anchor, GPI-anchor Apical cell membrane; Lipid-anchor,


GPI- anchor Basolateral cell membrane; Lipid-anchor, GPI-like-anchor. Secreted Cytoplasmic vesicle. Cytoplasmic vesicle, clathrin-coated vesicle. Endosome. Note=Endocytosed into cytoplasmic vesicles and then recycled to

the cell membrane


Tissue Location

Primarily expressed in tissues of epithelial origin. Expression is increased in malignant tissues. Expressed in kidney, lung and cerebellum. Detected in placenta and thymus epithelium.

Images



Anti-FOLR1 / FRA Reference Antibody (Dompe patent anti-FOLR1) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-FOLR1 / FRA Reference Antibody (Dompe patent anti-FOLR1)is more than 95% ,determined by SEC-HPLC.

Image not found: 202311/AP90912-4.jpg

Human GPC3 CHOS cells were stained with Anti-FOLR1 / FRA Reference Antibody (Dompe patent anti-FOLR1) and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC979=2.142 nM

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.