

# Anti-IL-5 Reference Antibody (Abgenix anti-IL-5)

Recombinant Antibody

Catalog # APR10958

## Product Information

---

<b>Application</b>	FC, Kinetics, Animal Model
<b>Primary Accession</b>	<a href="#">P05113</a>
<b>Reactivity</b>	Human, Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG4SP
<b>Calculated MW</b>	15238

## Additional Information

---

<b>Target/Specificity</b>	IL-5
<b>Endotoxin</b>	
<b>Conjugation</b>	Unconjugated
<b>Expression system</b>	CHO Cell
<b>Format</b>	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.


## Protein Information

---


<b>Name</b>	IL5
<b>Function</b>	Homodimeric cytokine expressed predominantly by T-lymphocytes and NK cells that plays an important role in the survival, differentiation, and chemotaxis of eosinophils (PubMed: <a href="#">2653458</a> , PubMed: <a href="#">9010276</a> ). Also acts on activated and resting B-cells to induce immunoglobulin production, growth, and differentiation (By similarity). Mechanistically, exerts its biological effects through a receptor composed of IL5RA subunit and the cytokine receptor common subunit beta/CSF2RB (PubMed: <a href="#">1495999</a> , PubMed: <a href="#">22528658</a> ). Binding to the receptor leads to activation of various kinases including LYN, SYK and JAK2 and thereby propagates signals through the RAS-MAPK and JAK-STAT5 pathways respectively (PubMed: <a href="#">7613138</a> ).
<b>Cellular Location</b>	Secreted.
<b>Tissue Location</b>	Present in peripheral blood mononuclear cells.

## Images

---



Anti-IL-5 Reference Antibody (Abgenix anti-IL-5) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-IL-5 Reference Antibody (Abgenix anti-IL-5) is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.