

Anti-MRC2 / CD280 Reference Antibody (Copenhagen Rigshospitalet patent anti-uPARAP)

Recombinant Antibody Catalog # APR10980

Product Information

Application FC, Kinetics, Animal Model

Primary Accession

Reactivity

Clonality

Isotype

Calculated MW

Q9UBG0

Human

Monoclonal

IgG1

166674

Additional Information

Target/Specificity MRC2 / CD280

Endotoxin

Conjugation Unconjugated

Expression system CHO Cell

Format Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

Protein Information

Name MRC2

Synonyms CLEC13E, ENDO180, KIAA0709, UPARAP

Function May play a role as endocytotic lectin receptor displaying calcium-dependent

lectin activity. Internalizes glycosylated ligands from the extracellular space for release in an endosomal compartment via clathrin-mediated endocytosis.

May be involved in plasminogen activation system controlling the

extracellular level of PLAUR/PLAU, and thus may regulate protease activity at

the cell surface. May contribute to cellular uptake, remodeling and

degradation of extracellular collagen matrices. May play a role during cancer progression as well as in other chronic tissue destructive diseases acting on collagen turnover. May participate in remodeling of extracellular matrix


cooperating with the matrix metalloproteinases (MMPs).

Cellular Location Membrane; Single-pass type I membrane protein.


Tissue Location Ubiquitous with low expression in brain, placenta, lung, kidney, pancreas,

spleen, thymus and colon. Expressed in endothelial cells, fibroblasts and

Images



Anti-MRC2 / CD280 Reference Antibody (Copenhagen Rigshospitalet patent anti-uPARAP) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-MRC2 / CD280 Reference Antibody (Copenhagen Rigshospitalet patent anti-uPARAP)is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.