

# Anti-Nogo Receptor / NgR Reference Antibody (Abbott patent anti-NGR)

Recombinant Antibody  
Catalog # APR10993

## Product Information

---

<b>Application</b>	FC, Kinetics, Animal Model
<b>Primary Accession</b>	<a href="#">Q9BZR6</a>
<b>Reactivity</b>	Human, Rat
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG2SA
<b>Calculated MW</b>	50708

## Additional Information

---

<b>Target/Specificity</b>	Nogo Receptor / NgR
<b>Endotoxin Conjugation</b>	Unconjugated
<b>Expression system</b>	CHO Cell
<b>Format</b>	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

## Protein Information

---

<b>Name</b>	RTN4R
<b>Synonyms</b>	NOGOR
<b>Function</b>	Receptor for RTN4, OMG and MAG (PubMed: <a href="#">12037567</a> , PubMed: <a href="#">12068310</a> , PubMed: <a href="#">12089450</a> , PubMed: <a href="#">12426574</a> , PubMed: <a href="#">12839991</a> , PubMed: <a href="#">16712417</a> , PubMed: <a href="#">18411262</a> , PubMed: <a href="#">19052207</a> ). Functions as a receptor for the sialylated gangliosides GT1b and GM1 (PubMed: <a href="#">18411262</a> ). Besides, functions as a receptor for chondroitin sulfate proteoglycans (By similarity). Can also bind heparin (By similarity). Intracellular signaling cascades are triggered via the coreceptor NGFR (PubMed: <a href="#">12426574</a> ). Signaling mediates activation of Rho and downstream reorganization of the actin cytoskeleton (PubMed: <a href="#">16712417</a> , PubMed: <a href="#">22325200</a> ). Mediates axonal growth inhibition (PubMed: <a href="#">12839991</a> , PubMed: <a href="#">19052207</a> , PubMed: <a href="#">28892071</a> ). Plays a role in regulating axon regeneration and neuronal plasticity in the adult central nervous system. Plays a role in postnatal brain development. Required for normal axon migration across the brain midline and normal formation of the corpus callosum. Protects motoneurons against apoptosis; protection against apoptosis is probably mediated via interaction

with MAG. Acts in conjunction with RTN4 and LINGO1 in regulating neuronal precursor cell motility during cortical development. Like other family members, plays a role in restricting the number dendritic spines and the number of synapses that are formed during brain development (PubMed:[22325200](https://pubmed.ncbi.nlm.nih.gov/22325200/)).


### Cellular Location

Cell membrane; Lipid- anchor, GPI-anchor. Membrane raft. Cell projection, dendrite {ECO:0000250|UniProtKB:Q99PI8}. Cell projection, axon {ECO:0000250|UniProtKB:Q99PI8}. Perikaryon {ECO:0000250|UniProtKB:Q99M75}. Note=Detected along dendrites and axons, close to synapses, but clearly excluded from synapses {ECO:0000250|UniProtKB:Q99PI8}


### Tissue Location

Widespread in the brain but highest levels in the gray matter. Low levels in heart and kidney; not expressed in oligodendrocytes (white matter).

## Images



Anti-Nogo Receptor / NgR Reference Antibody (Abbott patent anti-NGR) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-Nogo Receptor / NgR Reference Antibody (Abbott patent anti-NGR) is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.