

# Anti-TLR7 Reference Antibody (U.Tokyo patent anti-TLR7)

Recombinant Antibody

Catalog # APR11047

## Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	<a href="#">Q9NYK1</a>
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	120922

## Additional Information

Target/Specificity	TLR7
Endotoxin Conjugation	Unconjugated
Expression system	CHO Cell
Format	Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

## Protein Information

Name	TLR7 ( <a href="#">HGNC:15631</a> )
Function	<p>Endosomal receptor that plays a key role in innate and adaptive immunity (PubMed:<a href="#">14976261</a>, PubMed:<a href="#">32433612</a>). Controls host immune response against pathogens through recognition of uridine- containing single strand RNAs (ssRNAs) of viral origin or guanosine analogs (PubMed:<a href="#">12738885</a>, PubMed:<a href="#">27742543</a>, PubMed:<a href="#">31608988</a>, PubMed:<a href="#">32706371</a>, PubMed:<a href="#">35477763</a>). Upon binding to agonists, undergoes dimerization that brings TIR domains from the two molecules into direct contact, leading to the recruitment of TIR-containing downstream adapter MYD88 through homotypic interaction (PubMed:<a href="#">27742543</a>). In turn, the Myddosome signaling complex is formed involving IRAK4, IRAK1, TRAF6, TRAF3 leading to activation of downstream transcription factors NF-kappa-B and IRF7 to induce pro-inflammatory cytokines and interferons, respectively (PubMed:<a href="#">27742543</a>, PubMed:<a href="#">32706371</a>). In plasmacytoid dendritic cells, RNASET2 endonuclease cooperates with PLD3 or PLD4 5'-&gt;3' exonucleases to process RNA and release 2',3'-cyclic guanosine monophosphate (2',3'-cGMP) and cytidine-rich RNA fragments that occupy TLR7 ligand-binding pockets and trigger a signaling-competent state.</p> <p>Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:P58681};</p>


## Cellular Location

Single-pass type I membrane protein {ECO:0000250|UniProtKB:P58681}. Endosome {ECO:0000250|UniProtKB:P58681}. Lysosome {ECO:0000250|UniProtKB:P58681}. Cytoplasmic vesicle, phagosome {ECO:0000250|UniProtKB:P58681}. Note=Relocalizes from endoplasmic reticulum to endosome and lysosome upon stimulation with agonist {ECO:0000250|UniProtKB:P58681}


## Tissue Location

Detected in brain, placenta, spleen, stomach, small intestine, lung and in plasmacytoid pre-dendritic cells. Expressed in peripheral mononuclear blood cells (PubMed:32706371)

## Images



Anti-TLR7 Reference Antibody (U.Tokyo patent anti-TLR7) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-TLR7 Reference Antibody (U.Tokyo patent anti-TLR7) is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.