

Anti-LIV-1 / SLC39A6 Mouse Antibody

Catalog # APR11161

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	Q13433
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Calculated MW	85047

Additional Information

Target/Specificity	LIV-1 / SLC39A6
Expression system	CHO

Protein Information


Name	SLC39A6 (HGNC:18607)
Synonyms	LIV1, ZIP6
Function	Zinc-influx transporter which plays a role in zinc homeostasis and in the induction of epithelial-to-mesenchymal transition (EMT) (PubMed:12839489, PubMed:18272141, PubMed:21422171, PubMed:23919497, PubMed:27274087, PubMed:34394081). When associated with SLC39A10, the heterodimer formed by SLC39A10 and SLC39A6 mediates cellular zinc uptake to trigger cells to undergo epithelial- to- mesenchymal transition (EMT) (PubMed:27274087). The SLC39A10-SLC39A6 heterodimer also controls NCAM1 phosphorylation and its integration into focal adhesion complexes during EMT (By similarity). Zinc influx inactivates GSK3B, enabling unphosphorylated SNAI1 in the nucleus to down-regulate adherence genes such as CDH1, causing loss of cell adherence (PubMed:23919497). In addition, the SLC39A10-SLC39A6 heterodimer plays an essential role in initiating mitosis by importing zinc into cells to initiate a pathway resulting in the onset of mitosis (PubMed:32797246). Participates in the T-cell receptor signaling regulation by mediating cellular zinc uptake into activated lymphocytes (PubMed:21422171, PubMed:30552163, PubMed:34394081). Regulates the zinc influx necessary for proper meiotic progression to metaphase II (MII) that allows the oocyte-to-egg transition (PubMed:25143461).
Cellular Location	Cell membrane; Multi-pass membrane protein. Cell projection, lamellipodium membrane; Multi-pass membrane protein. Membrane raft; Multi-pass membrane protein. Apical cell membrane {ECO:0000250 UniProtKB:Q4V887} Note=Localizes to lipid rafts in T cells and is recruited into the immunological synapse in response to TCR stimulation (PubMed:34394081) In the choroid

plexus is limited to the apical membrane in epithelial cells (By similarity).
{ECO:0000250|UniProtKB:Q4V887, ECO:0000269|PubMed:34394081}


Tissue Location

Highly expressed in the breast, prostate, placenta, kidney, pituitary and corpus callosum (PubMed:12839489). Weakly expressed in heart and intestine. Also highly expressed in cells derived from an adenocarcinoma of the cervix and lung carcinoma (PubMed:12839489).


Images



Product on SDS-PAGE under non-reducing (NR) and reducing (R) conditions. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.



The purity of protein, determined by HPLC, is greater than 95%.



Human LIV1 CHOS cells were stained with anti-LIV-1 / SLC39A6 Mouse Antibody and negative control protein respectively, washed and then followed by PE and analyzed with FACS, EC50=0.2992 ug/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.