

Anti-IL-2Ra / CD25 Reference Antibody (Inolimomab)

Catalog # APR11195

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P01589
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	30819


Additional Information

Target/Specificity	IL-2Ra / CD25
Expression system	CHO

Protein Information



Name	IL2RA
Function	Receptor for interleukin-2. The receptor is involved in the regulation of immune tolerance by controlling regulatory T cells (TREGs) activity. TREGs suppress the activation and expansion of autoreactive T-cells.
Cellular Location	Membrane; Single-pass type I membrane protein.

Images



Anti-IL-2Ra / CD25 Reference Antibody (Inolimomab) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.

The purity of Anti-IL-2Ra / CD25 Reference Antibody (Inolimomab) is 99.5%, determined by SEC-HPLC.



Immobilized Human IL-2Ra / CD25, Fc Tag at 2 µg/mL
can bind Adintrevimab, EC50= 0.01263 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.