

Anti-B7-H1 / PD-L1 / CD274 Reference Antibody (Socazolimab)

Catalog # APR11204

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	Q9NZQ7
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	33275


Additional Information

Target/Specificity	B7-H1 / PD-L1 / CD274
Expression system	CHO


Protein Information

Name	CD274 (HGNC:17635)
Function	Plays a critical role in induction and maintenance of immune tolerance to self (PubMed: 11015443 , PubMed: 28813410 , PubMed: 28813417 , PubMed: 31399419). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed: 11015443 , PubMed: 28813410 , PubMed: 28813417 , PubMed: 36727298). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed: 10581077). Can also act as a transcription coactivator: in response to hypoxia, translocates into the nucleus via its interaction with phosphorylated STAT3 and promotes transcription of GSDMC, leading to pyroptosis (PubMed: 32929201).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Early endosome membrane; Single-pass type I membrane protein. Recycling endosome membrane; Single-pass type I membrane protein. Nucleus. Note=Associates with CMTM6 at recycling endosomes, where it is protected from being targeted for lysosomal degradation (PubMed: 28813417). Translocates to the nucleus in response to hypoxia via its interaction with phosphorylated STAT3 (PubMed: 32929201). [Isoform 2]: Endomembrane system; Single-pass type I membrane protein
Tissue Location	Highly expressed in the heart, skeletal muscle, placenta and lung. Weakly expressed in the thymus, spleen, kidney and liver. Expressed on activated T- and B-cells, dendritic cells, keratinocytes and monocytes.


Images



Anti-B7-H1 / PD-L1 / CD274 Reference Antibody (Socazolimab) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.



The purity of Anti-B7-H1 / PD-L1 / CD274 Reference Antibody (Socazolimab) is 98.5%, determined by SEC-HPLC.



Immobilized hu-PD-L1-ECD-His at 2 µg/mL can bind Socazolimab, EC50= 0.008638 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.