10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CTLA-4 / CD152 Reference Antibody (Tuvonralimab)

Catalog # APR11206

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P16410
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1-EM-DCCGKRE
Calculated MW	24656

Additional Information


Target/Specificity	CTLA-4 / CD152
Expression system	CHO


Protein Information

Name	CTLA4
Synonyms	CD152
Function	Inhibitory receptor acting as a major negative regulator of T-cell responses (PubMed:11279501, PubMed:11279502, PubMed:16551244, PubMed:1714933, PubMed:18641304, PubMed:28484017). Acts as a decoy receptor: the affinity of CTLA4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28 (PubMed:11279501, PubMed:11279502, PubMed:16551244, PubMed:1714933, PubMed:28484017).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Note=Exists primarily an intracellular antigen whose surface expression is tightly regulated by restricted trafficking to the cell surface and rapid internalization
Tissue Location	Widely expressed with highest levels in lymphoid tissues. Detected in activated T-cells where expression levels are 30- to 50-fold less than CD28, the stimulatory coreceptor, on the cell surface following activation.


Images

Anti-CTLA-4 / CD152 Reference Antibody (Tuvonralimab) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.





The purity of Anti-CTLA-4 / CD152 Reference Antibody (Tuvonralimab) is 99.14%, determined by SEC-HPLC.



Immobilized human CTLA4 Protein-His at 2 μg/mL can bind Tuvonralimab, EC50= 0.0100 μg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.