

Anti-SIRPa / CD172a Reference Antibody (Anzurstobart)

Catalog # APR11224

Product Information


Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P78324
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	54967

Additional Information


Target/Specificity	SIRPa / CD172a
Expression system	CHO

Protein Information


Name	SIRPA
Synonyms	BIT, MFR, MYD1, PTPNS1, SHPS1, SIRP
Function	Immunoglobulin-like cell surface receptor for CD47. Acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function (By similarity). Involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. Mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Plays a role in antiviral immunity and limits new world arenavirus infection by decreasing virus internalization (By similarity). Receptor for THBS1 (PubMed:24511121). Interaction with THBS1 stimulates phosphorylation of SIRPA (By similarity). In response to THBS1, involved in ROS signaling in non-phagocytic cells, stimulating NADPH oxidase-derived ROS production (PubMed:24511121).
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Ubiquitous. Highly expressed in brain. Detected on myeloid cells, but not T-cells. Detected at lower levels in heart, placenta, lung, testis, ovary, colon, liver, small intestine, prostate, spleen, kidney, skeletal muscle and pancreas



Anti-SIRPa / CD172a Reference Antibody (Anzurstobart) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.



The purity of Anti-SIRPa / CD172a Reference Antibody (Anzurstobart) is 98.2%, determined by SEC-HPLC.



Immobilized PHA247 Human SIRPA / CD172a, His Tag at 2 µg/mL can bind Anzurstobart, EC50= 0.004262 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.