

Anti-IL-13Ra1 Reference Antibody (Eblasakimab)

Catalog # APR11232

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P78552
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG4SP-opti
Calculated MW	48760

Additional Information



Target/Specificity	IL-13Ra1 / CD213a1
Expression system	CHO

Protein Information


Name	IL13RA1
Synonyms	IL13R, IL13RA
Function	Binds with low affinity to interleukin-13 (IL13). Together with IL4RA can form a functional receptor for IL13. Also serves as an alternate accessory protein to the common cytokine receptor gamma chain for interleukin-4 (IL4) signaling, but cannot replace the function of IL2RG in allowing enhanced interleukin-2 (IL2) binding activity.
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Ubiquitous. Highest levels in heart, liver, skeletal muscle and ovary; lowest levels in brain, lung and kidney Also found in B-cells, T-cells and endothelial cells

Images

Anti-IL-13Ra1 Reference Antibody (Eblasakimab) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.



The purity of Anti-IL-13Ra1 Reference Antibody (Eblasakimab) is 98.2%, determined by SEC-HPLC.



Immobilized IL-13R α 1-hFc at 2 μ g/mL can bind Eblasakimab, EC₅₀= 0.01244 μ g/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.