

Anti-TNFSF5 / CD40L / CD154 Reference Antibody (frexalimab)

Catalog # APR11245

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P29965
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1-ARA
Calculated MW	29274


Additional Information

Target/Specificity	TNFSF5 / CD40L / CD154
Expression system	CHO


Protein Information

Name	CD40LG
Synonyms	CD40L, TNFSF5, TRAP
Function	Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed: 1280226 , PubMed: 31331973). Costimulates T-cell proliferation and cytokine production (PubMed: 8617933). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed: 8617933). Induces the activation of NF-kappa-B (PubMed: 15067037 , PubMed: 31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed: 15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed: 15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).
Cellular Location	Cell membrane; Single-pass type II membrane protein. Cell surface
Tissue Location	Specifically expressed on activated CD4+ T- lymphocytes


Images



(frexalimab) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.



The purity of Anti-TNFSF5 / CD40L / CD154 Reference Antibody (frexalimab) is 99.3%, determined by SEC-HPLC.



PLVX-puro-TNFSF5(`CD40L)- FL-HEK293 cell line were stained with frexalimab and negative control protein respectively, washed and then followed by PE and analyzed with FACS, with the EC50 is 0.03 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.