

Anti-DLL3 Reference Antibody (FZ-A038)

Catalog # APR11278

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	Q9NYJ7
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1-LFLE-PS
Calculated MW	64618


Additional Information

Target/Specificity	DLL3
Expression system	CHO

Protein Information



Name	DLL3
Function	Inhibits primary neurogenesis. May be required to divert neurons along a specific differentiation pathway. Plays a role in the formation of somite boundaries during segmentation of the paraxial mesoderm (By similarity).
Cellular Location	Membrane; Single-pass type I membrane protein

Images



Anti-DLL3 Reference Antibody (FZ-A038) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.

The purity of Anti-DLL3 Reference Antibody (FZ-A038) is 99.22%, determined by SEC-HPLC.



Immobilized human-DLL3-ECD-His at 2 µg/mL can bind FZ-A038, EC50= 0.007668 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.