

Anti-STEAP2 Reference Antibody (Medimmune patent anti-STEAP2 CAR)

Catalog # APR11379

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	Q8NFT2
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	56056

Additional Information


Target/Specificity	STEAP2
Expression system	CHO

Protein Information


Name	STEAP2
Synonyms	PCANAP1, STAMP1
Function	Integral membrane protein that functions as a NADPH-dependent ferric-chelate reductase, using NADPH from one side of the membrane to reduce a Fe(3+) chelate that is bound on the other side of the membrane (By similarity). Mediates sequential transmembrane electron transfer from NADPH to FAD and onto heme, and finally to the Fe(3+) chelate (By similarity). Can also reduce Cu(2+) to Cu(1+) (By similarity).
Cellular Location	Endosome membrane {ECO:0000250 UniProtKB:Q8BWB6}; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein
Tissue Location	Expressed at high levels in prostate and at significantly lower levels in heart, brain, kidney, pancreas, and ovary.

Images

Anti-STEAP2 Reference Antibody (Medimmune patent anti-STEAP2 CAR) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.



The purity of Anti-STEAP2 Reference Antibody (Medimmune patent anti-STEAP2 CAR) is 100%, determined by SEC-HPLC.



PLVX-puro-STEAP2-FL cell line were stained with Medimmune patent anti-STEAP2 CAR and negative control protein respectively, washed and then followed by PE and analyzed with FACS, with the EC50 is 1.471 nM.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.