

Anti-PABPC1 Reference Antibody (alexstatug)

Catalog # APR11475

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P11940
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1-REM-nok
Calculated MW	70671

Additional Information

Target/Specificity	PABPC1
Expression system	CHO


Protein Information

Name	PABPC1 (HGNC:8554)
Function	Binds the poly(A) tail of mRNA, including that of its own transcript, and regulates processes of mRNA metabolism such as pre-mRNA splicing and mRNA stability (PubMed:11051545, PubMed:17212783, PubMed:25480299). Its function in translational initiation regulation can either be enhanced by PAIP1 or repressed by PAIP2 (PubMed:11051545, PubMed:20573744). Can probably bind to cytoplasmic RNA sequences other than poly(A) in vivo. Binds to N6-methyladenosine (m6A)-containing mRNAs and contributes to MYC stability by binding to m6A-containing MYC mRNAs (PubMed:32245947). Involved in translationally coupled mRNA turnover (PubMed:11051545). Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain (PubMed:11051545). Involved in regulation of nonsense-mediated decay (NMD) of mRNAs containing premature stop codons; for the recognition of premature termination codons (PTC) and initiation of NMD a competitive interaction between UPF1 and PABPC1 with the ribosome-bound release factors is proposed (PubMed:18447585). By binding to long poly(A) tails, may protect them from uridylation by ZCCHC6/ZCCHC11 and hence contribute to mRNA stability (PubMed:25480299).
Cellular Location	Cytoplasm. Cytoplasm, Stress granule. Nucleus. Cell projection, lamellipodium. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs (PubMed: 17289661). Shuttles between the cytoplasm and the nucleus (PubMed: 9582337). During stress and in the absence of DDX3X, localizes to the nucleus (PubMed: 21883093). At the leading edge of migrating fibroblasts, colocalizes with DDX3X (PubMed: 28733330). Relocalizes

to cytoplasmic stress granules upon cellular stress where it colocalizes with ENDOV (PubMed:27573237). In case of HRSV infection, localizes in cytoplasmic inclusion bodies substructures called inclusion bodies associated granules (IBAGs) (PubMed:31649314)

Tissue Location Ubiquitous.

Images



Anti-PABPC1 Reference Antibody (alextatug) on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 95%.

The purity of Anti-PABPC1 Reference Antibody (alextatug) is 98.71%, determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.