

FITC Anti-Human CD19 (4G7)

Catalog # APR11646

Product Information

Application	Functional assay, Kinetics (SPR), Kinetics (BLI), FACS, E, FTA
Primary Accession	P15391
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2b Fc silience
Calculated MW	61128


Additional Information

Target/Specificity	CD19
Expression system	CHO

Protein Information

Name	CD19
Function	Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes (PubMed: 29523808). Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed: 1373518 , PubMed: 16672701 , PubMed: 2463100). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed: 12387743 , PubMed: 16672701 , PubMed: 9317126 , PubMed: 9382888). Is not required for early steps during B cell differentiation in the blood marrow (PubMed: 9317126). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed: 1373518 , PubMed: 2463100). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed: 12387743 , PubMed: 16672701 , PubMed: 9317126).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Membrane raft {ECO:0000250 UniProtKB:P25918}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P25918}
Tissue Location	Detected on marginal zone and germinal center B cells in lymph nodes (PubMed: 2463100). Detected on blood B cells (at protein level) (PubMed: 16672701 , PubMed: 2463100)

Images



Human peripheral blood lymphocytes were stained with FITC Anti-Human CD19 (4G7) and APC anti-human CD3 (UCHT1), Analyzed data are derived from lymphocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.