

# NOD2 Antibody

Catalog # ASC10158

## Product Information

---

<b>Application</b>	WB, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q9HC29</a>
<b>Other Accession</b>	<a href="#">Q9HC29</a> , <a href="#">20137973</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	115283
<b>Concentration (mg/ml)</b>	1 mg/mL
<b>Conjugate</b>	Unconjugated
<b>Application Notes</b>	NOD2 antibody can be used for detection of Noxa by Western blot at 2 to 4 $\mu$ g/mL. Antibody can also be used for immunocytochemistry starting at 10 $\mu$ g/mL. For immunofluorescence start at 20 $\mu$ g/mL.

## Additional Information

---

<b>Gene ID</b>	64127
<b>Other Names</b>	NOD2 Antibody: CD, ACUG, BLAU, IBD1, NLRC2, NOD2B, CARD15, CLR16.3, PSORAS1, Nucleotide-binding oligomerization domain-containing protein 2, Caspase recruitment domain-containing protein 15, nucleotide-binding oligomerization domain containing 2
<b>Target/Specificity</b>	NOD2; NOD2 has no cross-reaction with NOD1.
<b>Reconstitution &amp; Storage</b>	NOD2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
<b>Precautions</b>	NOD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	NOD2 {ECO:0000303   PubMed:11087742, ECO:0000312   HGNC:HGNC:5331}
<b>Function</b>	Pattern recognition receptor (PRR) that detects bacterial peptidoglycan fragments and other danger signals and plays an important role in gastrointestinal immunity (PubMed: <a href="#">12514169</a> , PubMed: <a href="#">12527755</a> , PubMed: <a href="#">12626759</a> , PubMed: <a href="#">15044951</a> , PubMed: <a href="#">15998797</a> , PubMed: <a href="#">27283905</a> , PubMed: <a href="#">27748583</a> , PubMed: <a href="#">31649195</a> ). Specifically activated by muramyl dipeptide (MDP), a fragment of bacterial peptidoglycan found in every bacterial peptidoglycan type (PubMed: <a href="#">12514169</a> ,

PubMed:[12527755](#), PubMed:[12626759](#), PubMed:[12871942](#), PubMed:[15044951](#), PubMed:[15198989](#), PubMed:[15998797](#), PubMed:[22857257](#), PubMed:[23322906](#), PubMed:[27748583](#), PubMed:[36002575](#)). NOD2 specifically recognizes and binds 6-O-phospho-MDP, the phosphorylated form of MDP, which is generated by NAGK (PubMed:[36002575](#)). 6-O-phospho-MDP-binding triggers oligomerization that facilitates the binding and subsequent activation of the proximal adapter receptor-interacting RIPK2 (PubMed:[11087742](#), PubMed:[17355968](#), PubMed:[21887730](#), PubMed:[23806334](#), PubMed:[28436939](#)). Following recruitment, RIPK2 undergoes 'Met-1'- (linear) and 'Lys-63'-linked polyubiquitination by E3 ubiquitin-protein ligases XIAP, BIRC2, BIRC3 and the LUBAC complex, becoming a scaffolding protein for downstream effectors, triggering activation of the NF-kappa-B and MAP kinases signaling (PubMed:[11087742](#), PubMed:[12514169](#), PubMed:[12626759](#), PubMed:[15198989](#), PubMed:[21887730](#), PubMed:[23322906](#), PubMed:[23806334](#), PubMed:[28436939](#)). This in turn leads to the transcriptional activation of hundreds of genes involved in immune response (PubMed:[15198989](#)). Its ability to detect bacterial MDP plays a central role in maintaining the equilibrium between intestinal microbiota and host immune responses to control inflammation (By similarity). An imbalance in this relationship results in dysbiosis, whereby pathogenic bacteria prevail on commensals, causing damage in the intestinal epithelial barrier as well as allowing bacterial invasion and inflammation (By similarity). Acts as a regulator of appetite by sensing MDP in a subset of brain neurons: microbiota-derived MDP reach the brain, where they bind and activate NOD2 in inhibitory hypothalamic neurons, decreasing neuronal activity, thereby regulating satiety and body temperature (By similarity). NOD2- dependent MDP-sensing of bacterial cell walls in the intestinal epithelial compartment contributes to sustained postnatal growth upon undernutrition (By similarity). Also plays a role in antiviral response by acting as a sensor of single-stranded RNA (ssRNA) from viruses: upon ssRNA-binding, interacts with MAVS, leading to activation of interferon regulatory factor-3/IRF3 and expression of type I interferon (PubMed:[19701189](#)). Also acts as a regulator of autophagy in dendritic cells via its interaction with ATG16L1, possibly by recruiting ATG16L1 at the site of bacterial entry (PubMed:[20637199](#)). NOD2 activation in the small intestine crypt also contributes to intestinal stem cells survival and function: acts by promoting mitophagy via its association with ATG16L1 (By similarity). In addition to its main role in innate immunity, also regulates the adaptive immune system by acting as regulator of helper T-cell and regulatory T-cells (Tregs) (By similarity). Besides recognizing pathogens, also involved in the endoplasmic reticulum stress response: acts by sensing and binding to the cytosolic metabolite sphingosine-1-phosphate generated in response to endoplasmic reticulum stress, initiating an inflammation process that leads to activation of the NF-kappa-B and MAP kinases signaling (PubMed:[27007849](#), PubMed:[33942347](#)). May also be involved in NLRP1 activation following activation by MDP, leading to CASP1 activation and IL1B release in macrophages (PubMed:[18511561](#)).

#### Cellular Location

Cell membrane; Lipid-anchor. Basolateral cell membrane. Cytoplasm Mitochondrion. Note=Palmitoylation promotes localization to the cell membrane, where it detects bacterial invasion at the point of entry.

#### Tissue Location

Expressed in monocytes, macrophages, dendritic cells, hepatocytes, preadipocytes, epithelial cells of oral cavity, lung and intestine, with higher expression in ileal Paneth cells and in intestinal stem cells.

## Background

NOD2 Antibody: Apaf-1 and NOD1 are members of a new family, which are involved in the regulation of

apoptosis and immune response. Each of them contains a caspase recruitment domain (CARD) and a nucleotide-binding oligomerization domain (NOD). A third member in this family was recently identified and designated NOD2. NOD2 interacts with RICK via a homophilic CARD-CARD interaction. NOD2 activates NF- $\kappa$ B, which is regulated by its carboxy-terminal leucine-rich repeat domain that acts as an intracellular receptor for components of bacteria. The variants of NOD2, either a frameshift or a missense, were associated with Crohn's disease that is a main type of chronic inflammatory bowel disease.


## References

---


- Inohara N, Koseki T, del Peso L, et al. Nod1, an Apaf-1-like activator of caspase-9 and nuclear factor- $\kappa$ B. *J. Biol. Chem.* 1999; 274:14560-7.
- Ogura Y, Inohara N, Benito A, et al. Nod2, a Nod1/Apaf-1 family member that is restricted to monocytes and activates NF- $\kappa$ B. *J. Biol. Chem.* 2001; 276:4812-8.
- Hugot JP, Chamaillard M, et al. Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* 2001; 411:599-603.
- Ogura Y, Bonen DK, Inohara N, et al. A frameshift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* 2001; 411:603-6.

## Images


---



Western blot analysis of NOD2 in HeLa cell lysate with NOD2 antibody at (A) 2 and (B) 4  $\mu$ g/mL.



Immunocytochemistry of NOD2 in HeLa cells with NOD2 antibody at 10  $\mu$ g/mL.



Immunofluorescence of NOD2 in HeLa cells with NOD2 antibody at 20  $\mu$ g/mL.