

Lano Antibody

Catalog # ASC10161

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	Q9BTT6
Other Accession	AAK72246 , 14701834
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	59242
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	Lano antibody can be used for detection of Lano by Western blot at 1 - 2 μ g/mL. Antibody can also be used for immunohistochemistry starting at 10 μ g/mL. For immunofluorescence start at 20 μ g/mL.

Additional Information

Gene ID	55227
Other Names	Lano Antibody: LANO, dj523E19.1, LANO, Leucine-rich repeat-containing protein 1, LANO adapter protein, leucine rich repeat containing 1
Target/Specificity	LRRC1;
Reconstitution & Storage	Lano antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	Lano Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	LRRC1
Synonyms	LANO
Cellular Location	Cytoplasm. Membrane; Peripheral membrane protein. Note=Localized at the basolateral side of epithelial cells
Tissue Location	Expressed strongly in testis and placenta, followed by heart, lung, kidney, thyroid, trachea, colon, prostate and pancreas


Background

Lano Antibody: Lano is a member of the LAP (leucine-rich repeats and PDZ) family of proteins that also includes Densin-180, Erbin, and hScribble. The LAP proteins generally contain multiple leucine-rich repeat (LRR) domains which serve to target them to the basolateral membrane of epithelial cells. Lano is unique in that it alone does not possess one or more PDZ (PSD95/DLG/ZO-1) domains as do the other members of the LAP family. However, it can bind to the PDZ domain of Erbin in addition to those of membrane-associated and guanylate kinase (MAGUK) proteins which regulate adhesion and plasticity at cell junctions. It has been suggested that it is through these interaction that these LAP proteins participate in the maintenance of proper embryonic development and integrity of epithelial tissues.


References

Saito H, Santoni M-J, Jaulin-Bastard F, et al. Lano, a novel LAP protein directly connected to MAGUK proteins in epithelial cells. *J. Biol. Chem.* 2001; 276:32051-5.
Legouis R, Jaulin-Bastard F, Schott S, et al. Basolateral targeting by leucine-rich repeat domains in epithelial cells. *EMBO Rep.* 2003; 4:1096-102.
Funke L, Dakoji S, Bredt DS. Membrane-Associated Guanylate Kinases Regulate Adhesion and Plasticity at Cell Junctions. *Annu. Rev. Biochem.* 2005; 74:219-45.
Bilder D and Perrimon N. Localization of apical epithelial determinants by the basolateral PDZ protein Scribble. *Nature* 2000; 403:676-80.


Images



Western blot analysis of Lano in PC-3 whole cell lysate with Lano antibody at (A) 1 or (B) 2 $\mu\text{g}/\text{ml}$.



Immunohistochemistry of Lano in human colon tissue with Lano antibody at 10 $\mu\text{g}/\text{mL}$.



Immunofluorescence of Lano in human colon tissue with Lano antibody at 20 $\mu\text{g}/\text{mL}$.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.