

# NALP2 Antibody

Catalog # ASC10169

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q9NX02</a>
<b>Other Accession</b>	<a href="#">NP_060322</a> , <a href="#">8923473</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	120515
<b>Concentration (mg/ml)</b>	1 mg/mL
<b>Conjugate</b>	Unconjugated
<b>Application Notes</b>	NALP2 antibody can be used for detection of NALP2 by Western blot at 1 to 2 $\mu$ g/mL.

## Additional Information

---

<b>Gene ID</b>	55655
<b>Other Names</b>	NALP2 Antibody: NBS1, PAN1, NALP2, PYPAF2, CLR19.9, NBS1, NACHT, LRR and PYD domains-containing protein 2, Nucleotide-binding site protein 1, NLR family, pyrin domain containing 2
<b>Target/Specificity</b>	NLRP2;
<b>Reconstitution &amp; Storage</b>	NALP2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
<b>Precautions</b>	NALP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	NLRP2
<b>Synonyms</b>	NALP2, NBS1, PAN1, PYPAF2
<b>Function</b>	Suppresses TNF- and CD40-induced NFKB1 activity at the level of the IKK complex, by inhibiting NFKBIA degradation induced by TNF. When associated with PYCARD, activates CASP1, leading to the secretion of mature pro-inflammatory cytokine IL1B. May be a component of the inflammasome, a protein complex which also includes PYCARD, CARD8 and CASP1 and whose function would be the activation of pro-inflammatory caspases.

**Cellular Location**

Cytoplasm

**Tissue Location**

Expressed at high levels in lung, placenta and thymus and at lower levels in ovary, intestine and brain (PubMed:15456791). Highly abundant in oocytes and early embryos, however poorly expressed in somatic tissues such as brain, kidney, liver and spinal cord (PubMed:30877238).


## Background

**NALP2 Antibody:** NALP2 belongs to a family of cytoplasmic proteins that have been implicated in cell responses to apoptotic and inflammatory stimuli. Unlike the prototypical NALP protein NALP1, NALP2 only contains a NACHT domain, leucine rich repeat (LRR), and pyrin-containing domain (PYD). This protein interacts with the adapter protein ASC in addition to CARD8 and caspase-1 to form an inflammasome with high proIL-1 $\beta$ -processing activity and is thought to function as a modulator of NF- $\kappa$ B and procaspase-1 activation in macrophages. It has also been suggested that NALP2, in addition to other NALP family members, can act as innate sensors of pathogens in a manner similar to the toll-like receptors (TLRs). At least two alternatively spliced transcript variants encoding distinct isoforms have been found for NALP2.

## References

- Tschopp J, Martinon F, and Burns K. NALPs: a novel protein family involved in inflammation. *Nat. Rev. Mol. Cell Biol.* 2003; 4:95-104.
- Bruey JM, Bruey-Sedano N, Newman R, et al. PAN1/NALP2/PYPAF2, an inducible inflammatory mediator that regulates NF- $\kappa$ B and caspase-1 activation in macrophages. *J. Biol. Chem.* 2004; 279:51897-907.
- Agostini L, Martinon F, Burns K, et al. NALP3 forms an IL-1 $\beta$ -processing inflammasome with increased activity in Muckle-Wells autoinflammatory disorder. *Immunity* 2004; 20:319-25.
- Martinon F and Tschopp J. NLRs join TLRs as innate sensors of pathogens. *TRENDS Imm.* 2005; 26:447-54.

## Images



Western blot analysis of NALP2 in PC-3 cell lysate with NALP2 antibody at (A) 1 and (B) 2  $\mu$ g/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.